

Exercise 2: Grover's algorithm and lower bounds


Lecturer: Simon Apers (apers@irif.fr)

Exercise 1 (Amplitude amplification). A useful variation on Grover's algorithm is called *amplitude amplification*. Assume that we have access to a unitary U (and its inverse U^\dagger) such that


$$U |0^n\rangle |0\rangle = |\psi\rangle = \sqrt{p} |\psi_1\rangle |1\rangle + \sqrt{1-p} |\psi_0\rangle |0\rangle,$$

and we would like to prepare the "marked" state $|\psi_1\rangle$.


- The following circuit presents a simple solution. What is its success probability?



Amplitude amplification improves on this. Consider the amplitude amplification operator:



- For states of the form $\alpha |\psi_1\rangle |1\rangle + \beta |\psi_0\rangle |0\rangle$, show that this circuit corresponds to a product of a reflection around $|\psi_0\rangle |0\rangle$ and a reflection around $|\psi\rangle$.
- What is the success probability of the following circuit?



Exercise 2 (Multilinear polynomials). Show that any function $f : \{0, 1\}^N \rightarrow \mathbb{C}$ has a unique representation as a multilinear polynomial of degree at most N .

Exercise 3 (Symmetric functions). A function $\{0, 1\}^N \rightarrow \mathbb{C}$ is called a *symmetric* function if $f(x)$ only depends on the Hamming weight $|x|$ (i.e., there exists a function \bar{f} such that $f(x) = \bar{f}(|x|)$). Examples are the OR-function and the PARITY-function. Through a symmetrization argument, one can show that $\deg(f) = \deg(\bar{f})$ and $\widetilde{\deg}(f) = \widetilde{\deg}(\bar{f})$.

- If f is the PARITY-function on N bits, show that $\deg(\bar{f}) \geq N$. This implies that any zero-error quantum algorithm for PARITY must make $N/2$ quantum queries.
- If f is the PARITY-function on N bits, show that $\widetilde{\deg}(\bar{f}) \geq N$. This implies that even a bounded-error quantum algorithm for PARITY must make $N/2$ quantum queries.

Exercise 4 (Quantum approximate counting). Check that the Grover operator G has eigenvectors and corresponding eigenvalues

$$|\psi_{\pm}\rangle = \frac{|u_1\rangle \pm i|u_0\rangle}{\sqrt{2}}, \quad \lambda_{\pm} = e^{\pm 2i\theta}.$$

Use quantum phase estimation on the initial state

$$|u\rangle = \frac{-i}{\sqrt{2}}(e^{i\theta}|\psi_+\rangle - e^{-i\theta}|\psi_-\rangle).$$

to estimate θ (and hence t/N).